

TRINITY COLLEGE DUBLIN
School of Computer Science and Statistics

Extra Questions

ST3009: Statistical Methods for Computer Science

Question 1. Suppose two continuous valued random variables X and Y have the following joint PDF

$$f_{XY}(x, y) = \begin{cases} 0 & x < 0, y < 0 \\ 1 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & x > 1, y > 1 \end{cases}$$




Figure 1: Plot of PDF $f_{XY}(x, y)$

- (a) Calculate $P(0 \leq X \leq 0.5 \text{ and } 0 \leq Y \leq 0.5)$?
 (b) Calculate $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5)$?

Solution

- $P(0 \leq X \leq 0.5 \text{ and } 0 \leq Y \leq 0.5)$ is the volume under the PDF for $0 \leq X \leq 0.5$ and $0 \leq Y \leq 0.5$. This is a cube of length 0.5, breadth 0.5 and height 1, so its volume is $0.5 \times 0.5 \times 1 = 0.25$.
- We split the volume under the PDF for $0 \leq X \leq 2$ and $0 \leq Y \leq 0.5$ into two parts, the volume under the PDF for $0 \leq X \leq 1$ and $0 \leq Y \leq 0.5$ and the volume $1 \leq X \leq 2$ and $0 \leq Y \leq 0.5$. The volume under the PDF for $0 \leq X \leq 1$ and $0 \leq Y \leq 0.5$ is a cube of length 1, breadth 0.5 and height 1 so its volume is $1 \times 0.5 \times 1 = 0.5$. The volume under the PDF for $1 \leq X \leq 2$ and $0 \leq Y \leq 0.5$ is 0. So the total volume is 0.5 i.e. $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5) = 0.5$

Question 2. Suppose two continuous valued random variables X and Y have the following joint CDF

$$F(x, y) = \begin{cases} 0 & x < 0, y < 0 \\ xy & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ y, x > 1, 0 \leq y \leq 1 \\ x, y > 1, 0 \leq x \leq 1 \\ 1 & x > 1, y > 1 \end{cases}$$

- (a) Sketch the graph of this CDF.
- (b) Calculate $P(X \leq 0.5 \text{ and } Y \leq 0.5)$
- (c) Calculate $P(0.1 \leq X \leq 0.5 \text{ and } 0.1 \leq Y \leq 0.5)$?
- (d) Calculate $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5)$?
- (e) Are X and Y independent ? Hint: recall $P(X \leq x) = F(x, \infty)$.

Solution

- $P(X \leq 0.5 \text{ and } Y \leq 0.5) = F(0.5, 0.5) = 0.25$
- $P(0 \leq X \leq 0.5 \text{ and } 0 \leq Y \leq 0.5) = F(0.5, 0.5) - F(0.1, 0.1) = 0.25 - 0.01 = 0.024$
- $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5) = F(2, 0.5) - F(0, 0) = 0.5 - 0 = 0.5$
- For independence we need $P(X \leq x \text{ and } Y \leq y) = P(X \leq x)P(Y \leq y)$. Now $P(X \leq x) = F(x, \infty) = x$ for $0 \leq x \leq 1$ and $P(Y \leq y) = F(\infty, y) = y$ for $0 \leq y \leq 1$, so $P(X \leq x)P(Y \leq y) = xy$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$. But $P(X \leq x \text{ and } Y \leq y) = F(x, y) = xy$ when $0 \leq x \leq 1$ and $0 \leq y \leq 1$. That is, $P(X \leq x \text{ and } Y \leq y) = P(X \leq x)P(Y \leq y)$. We can repeat for $x < 0$ and $x > 1$ etc to confirm that this holds for all values of x and y . Hence, X and Y are independent.

Question 3. Suppose two continuous valued random variables X and Y have the following joint CDF

$$F(x, y) = \begin{cases} 0 & x < 0, y < 0 \\ x^2y/2 + xy^3/2 & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ x^2/2 + x/2 & 0 \leq 0 \leq x \leq 1, y \geq 1 \\ y/2 + y^3/2, x > 1, 0 \leq y \leq 1 \\ 1 & x > 1, y > 1 \end{cases}$$

- (a) Calculate $P(X \leq 0.5 \text{ and } Y \leq 0.5)$
- (b) Calculate $P(0.1 \leq X \leq 0.5 \text{ and } 0.1 \leq Y \leq 0.5)$?
- (c) Calculate $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5)$?

Solution

- $P(X \leq 0.5 \text{ and } Y \leq 0.5) = F(0.5, 0.5) = 0.5^2 \times 0.5/2 + 0.5 \times 0.5^3/2 = 0.0938$
- $P(0 \leq X \leq 0.5 \text{ and } 0 \leq Y \leq 0.5) = F(0.5, 0.5) - F(0.1, 0.1) = 0.0938 - 0.00055 = 0.0932$
- $P(0 \leq X \leq 2 \text{ and } 0 \leq Y \leq 0.5) = F(2, 0.5) - F(0, 0) = 0.5/2 + 0.5^3/2 - 0 = 0.3125$

Question 4. Suppose random variables X and Y have PDFs $f_X(x) = e^{-x}$, $f_Y(y) = 0.5e^{-0.5y}$. Suppose also the X and Y are independent.

- (a) What is their joint PDF ?
- (b) Sketch a graph of this PDF.

Solution

- Since they are independent we just multiply the individual PDFs. That is, $f_{XY}(x, y) = 0.5e^{-x}e^{-0.5y} = 0.5e^{-x-0.5y}$




Figure 2: Plot of PDF $f_{XY}(x, y)$

Question 5. Suppose random variable X had PDF

$$f_X(x) = \begin{cases} 0 & x \leq 0 \\ 1 & 0 < x \leq 1 \\ 0 & x > 1 \end{cases}$$

and random variable Y has PDF

$$f_Y(y) = \begin{cases} 0 & y \leq 0.5 \\ 1 & 0.5 < y \leq 1.5 \\ 0 & y > 1.5 \end{cases}$$

Suppose also the X and Y are independent.

- (a) What is their joint PDF ?
- (b) Sketch a graph of this PDF.

Solution

- Since they are independent we just multiply the individual PDFs. That is,

$$f_{XY}(x, y) = \begin{cases} 0 & x \leq 0 \text{ or } y \leq 0.5 \\ 1 & 0 < x \leq 1, 0.5 < y \leq 1.5 \\ 0 & x > 1 \text{ or } y > 1.5 \end{cases}$$

Question 6. Suppose two random variables X and Y have PDFs $f_X(x) = e^{-x}$, $f_Y(y) = 0.5e^{-0.5y}$ and conditional PDF $f_{Y|X}(y|x) = e^{-|x-y|}$. Using Bayes Rule for PDFs write an expression for $f_{X|Y}(x|y)$.

Solution

- $f_{X|Y}(x|y) = f_{Y|X}(y|x)f_X(x)/f_Y(y) = e^{-|x-y|}e^{-x}/(0.5e^{-y}) = 2e^{-|x-y|-x+y}$




Figure 3: Plot of PDF $f_{XY}(x,y)$